

<https://helda.helsinki.fi>

---

## Vakava puudutemyrkytys

Rosenberg, Per

2019

---

Rosenberg , P 2019 , ' Vakava puudutemyrkytys ' , Finnanest. , Vuosikerta. 52 , Nro 1 , Sivut 22-23 . < [http://www.finnanest.fi/files/rosenberg\\_vakava\\_puudutemyrkytys.pdf](http://www.finnanest.fi/files/rosenberg_vakava_puudutemyrkytys.pdf) >

---

<http://hdl.handle.net/10138/302281>

---

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



**Per Rosenberg**

Helsingin yliopisto,  
anestesiologian professori (emer.)  
per.rosenberg@fimnet.fi

# Vakava puudutemyrkytys

**Suomessa anestesiologien suorittamiin puudutuksiin liittyvät vakavat puudutemyrkytykset ovat onneksi harvinaisia (1). Itse olen urallani kohdannut muutaman, joista tässä selostan vakavimman. Vuosi oli 1982.**

► Yksittäiset tapauselostukset johtavat harvoin klinisen käytännön muutoksiin, mutta tämä tapaus ja sen syyn selvittäminen herättivät huomiota myös kansainvälisesti vaikuttaen osaltaan siihen, että bupivakaiinin käyttö laskimoanestesiassa (IVRA) loppui tai väheni huomattavasti. Potilaan selviämiseen ja hyvään kuntoon toipumiseen saattoi vaikuttaa se, että myrkytystä hoitamassa oli kaksi anestesiologian erikoislääkärinä. Koska aikaa tästä potilastapauksesta on kulunut paljon

## IVRA-puudutuksessa osa puuduteannoksesta voi päästä yleiseen verenkiertoon.

eikä muistiin voi aina luottaa, olen mahdollisimman tarkasti poiminut oleelliset yksityiskohdat siitä amerikkalaisen lehden julkaisusta, jossa tapauselostus julkaistiin vuonna 1983 (2) (kuva 1). Käsillä olevassa artikkelissa olen pyrkinyt kursivoidulla tekstillä tekemään pieniä selvennyksiä alkuperäisiin tietoihin.

### Potilastapaus

Potilaana oli 71-vuotias mies (120 kg, 180 cm), joka tuli ekskisiroleikkaukseen Dupuytrenin kutistuskoukistusta

man vuoksi. Eteisvärinän vuoksi hän käytti digoksiinia ja pindololia. Verenpaine oli 160/80 mmHg leikkausta edeltävänä päivänä. Esilääkkeeksi potilas sai 20 mg diatsepaamia suun kautta ja 80 mg petidiiniä lihakseen noin 60 minuuttia ennen anestesian alkua.

Raajan verityhjiö aikaansaatiin Esmarchin siteellä, ja proksimaalinen kahdesta erillisestä painemansetista (Kidde and Co, Warsaw, Indiana) täytettiin 300 mmHg:n paineeseen. Olkavarren ympärys m. bicepsin kohdalla oli 42–44 cm (mitattuna kaksi päivää myöhemmin) ja mansetti oli 7,5 cm leveä ja 65 cm pitkä. Esmarchin siteen poiston jälkeen laskimoanestesiaa varten ruiskutettiin 80 ml 0,25 %:sta bupivakaiinia perifeeriseen laskimoon noin 30 sekunnissa. Tällöin verenpaine mitattuna vastakkaisesta yläraajasta oli 170–180/110 mmHg.

Noin kaksi minuuttia bupivakaiinin ruiskuttamisen jälkeen potilas meni äkisti bradykardiseksi ja synoottiseksi, ja samalla tajuttomaksi (potilas kuvasi jälkeensä tunteen vastanneen yleisanestesian alkua). Muutama sekunti näiden ensioireiden jälkeen alkoivat kouristukset, jotka voimistuivat *grand mal*-kouristuksiksi. Verenpainetta ei pystytty mittaamaan. *Kouristukset häiritsivät EKG-käyrää, joten epäselväksi jäi mikä oli sydämen*

*rytmi näiden kriittisten vaiheiden aikana.* Käsin ventiloiminen (FIO<sub>2</sub> 1,0) onnistui, kun laskimoon oli annettu tiopentaalia 150 mg ja suksametonia 150 mg. Potilaan henkitorvi intuboitui ja painelupuhalluselvytys aloitettiin. Noin 15 minuuttia kestäneen elvytyksen aikana potilaalle annettiin adrenaliinia 0,5 mg kahdesti, kalsiumkloridia 1,5 g sekä atropiinia 1 mg i.v. Natriumbikarbonaatin kokonaismäärä oli 270 mekv perustuen valtimoverikaasumäärittelyyn 10 minuutin kuluttua elvytyksen alusta (pH 7,09, PCO<sub>2</sub> 56 mmHg, PO<sub>2</sub> 160 mmHg, BE –14,4 mekv/l) ja 25 minuutin kuluttua elvytyksen alusta (pH 7,19, PCO<sub>2</sub> 45,8 mmHg, PO<sub>2</sub> 106,8 mmHg, BE –11,1 mekv/l). Spontaaniverenkierron palautumisen jälkeen hemodynaamiikkaa tuettiin dopamiini- ja nitroprussidi-laskimoinfuusioilla.

Potilaan tajunta palasi noin 30 minuutin kuluttua elvytyksen aloituksesta ja verenkierron tila palasi preoperatiivista vastaavaksi (EKG:ssa eteisvärinä). Bupivakaiinin valtimoveripitoisuudet 10 minuuttia, 25 minuuttia ja 40 minuuttia myrkytysoireiden ilmaantumisen alusta olivat 1,8 µg/ml, 0,9 µg/ml ja 0,58 µg/ml.

Leikkausta ei tehty. Verityhjiömansetti pidettiin täytettynä 300 mmHg:n paineella yhteensä 55 minuuttia, ja ennen paineen poistamista potilaalle annettiin tiopen-

taalia 100 mg. Uusia myrkytysoireita ei ilmaantunut ja leikattavan puolen solislaskimoveren bupivakaiinipitoisuudet yhden ja viiden minuutin kulluttua olivat 1,05 µg/ml ja 0,65 µg/ml.

Potilas toipui hyvin eikä kliinisiä merkkejä hypoksisista elinvaurioista voitu todeta.

## Täydentävä ja selventävä tutkimus

Kaksi päivää myöhemmin potilaalle tehtiin flebografia hoidettavaan raajaan käyttäen samaa tekniikkaa, välineistöä ja injektaattilavuutta kuin myrkytykseen johtaneessa IVRA-puudutuksessa. Röntgenkuvis- näkyi selkeästi, että paineistetusta mansetista huolimatta osa varjoaineesta tunkeutui solislaskimoon.

Potilaan verenpaine flebografian aikana oli 170–190/100–110 mmHg. Jälkeenpäin, ja täydennyksenä tälle alkuperäiselle tapausselostukselle, tehtiin kuudelle vapaaehtoiselle koehenkilölle flebografiatutkimus IVRA-olosuhteita simuloiden. Aineisto koostui kolmesta aikaisemmin

hoidetusta käsileikkauspotilaasta, joilla IVRA-puudutuksen aikana oli havaittu mitattavia puudutepitoisuuksia verenkierrassa verityhjiömansetin ollessa täytettynä, sekä kolmesta normaalkokoisesta terveestä vapaaehtoisesta henkilöstä. Kokeen aikana verityhjiömansetin paine oli 250 mmHg neljällä henkilöllä ja 300 mmHg kahdella. Yhteensä neljällä henkilöllä (kahdella kummastakin ryhmästä) näkyi pieniä määriä varjoainetta joko mansetin proksimaalipuolella tai viiruina mansetin alla.

## Johdopäätökset

IVRA-puudutuksessa osa puuduteannoksesta voi päästä yleiseen verenkiertoon täytetystä olkavarsimansetista huolimatta. On oletettavaa, että puudutteen läpipääsyriski täytetyn mansetin alta kasvaa, jos potilas on lihaksikas ja valtimoverenpaine on korkea (kuten selostetussa potilastapauksessa).

Asidoosi pahentaa puudute-myrkytystä. Hyvästä ventilaatiosta ja natriumbikarbonaatin annosta huo-

limatta myrkytyspotilaamme ajautui metaboliseen asidoosiin. Vakavissa myrkytyksissä valtimoverikaasujen vierimäärityksillä on tärkeä hoitoa ohjaava merkitys.

Vakavassa puudutemyrkytyksessä kouristukset haittaavat sydämen toiminnan monitorointia ja verenpaineen mittausta. Ne vaikeuttavat myös ventilaatiota ja intubaatiota. Diatsepaami on tehokas lievän keskushermostoärsytyksen hoidossa, mutta voimakkaat ja elvytystoimenpiteitä vaativat kouristukset hoituvat parhaiten laskimonsisäisellä anestetiilla (nykyään propofolilla). ■

## Viitteet

1. Heinonen JA, Litoinen E, Pitkänen M, Rosenberg PH. Incidence of severe local anaesthetic toxicity and adoption of lipid rescue in Finnish anaesthesia departments in 2011–2013. *Acta Anaesthesiol Scand* 2015; 59: 1032–1037.
2. Rosenberg PH, Kalso EA, Tuominen MK, Lindén HB. Acute bupivacaine toxicity as a result of venous leakage under the tourniquet cuff during a Bier block. *Anesthesiology* 1983; 58: 95–98.

CLINICAL REPORTS 95

Anesthesiology  
1983;58:95, 1993

### Acute Bupivacaine Toxicity As a Result of Venous Leakage Under the Tourniquet Cuff during a Bier Block

PER H. ROSENBERG, M.D.,\* EJA A. KALSO, M.D.,† MARJATTA E. TUOMINEN, M.D.,‡  
HANS B. LINDÉN, M.D.§

Severe cardiovascular complications have been associated with regional anesthesia using the newer local anesthetics at clinical<sup>1</sup> or larger doses.<sup>2</sup> Unintentional intravascular injection or rapid absorption has been considered responsible for these reactions. Lately, bupivacaine has become increasingly popular for the Bier block because it produces residual analgesia after the tourniquet is released.<sup>3,4</sup> A dose as high as 60 ml of 0.5% bupivacaine has been recommended,<sup>5</sup> although the use of 40–60 ml of 0.25% bupivacaine in adults has been considered sufficient and safe.<sup>6</sup> We have routinely used a 0.25% solution and a dose of approximately 1.8–2.0 mg/kg with satisfactory blocking results.<sup>7</sup>

Recently, we encountered a case of bupivacaine intoxication soon after the injection of the anesthetic. Because the tourniquet cuff appeared intact, we performed a tourniquet phlebography of the arm in this patient and six other subjects to see if local anesthetic was entering the general circulation.

#### REPORT OF A CASE

A 71-year-old man (170 kg, 180 cm) was scheduled for excision of a Desmoplasty carcinoma. He was taking digoxin and glyceryl trinitrate before surgery. Diuretics, 20 mg po, and aspirin, 80 mg po, were given about 60 min prior to anesthesia. The arm was exsanguinated with an Esmarch bandage and the proximal of the two separate pneumatic cuffs (Käthe and Co., Witten, Ireland) was inflated to 200 mmHg. The circumference of the arm at the biceps region was about 43–44 cm, and the cuff was 7.8 cm wide and 48 cm long. After removal of the Esmarch bandage, 100 ml of 0.25% bupivacaine (IVRA) was performed by injecting 60 ml of 0.25% bupivacaine into a vein of the wrist in about 30 s. Arterial blood pressure recorded in the contralateral arm was 170/100/110 mmHg. After about two minutes after the injection of bupivacaine, the patient suddenly developed tachycardia and cyanosis, and lost consciousness (pupils were reacted only to a feeling of having received general anesthesia).

#### STUDY SUBJECTS AND METHODS

In a previous study (performed about eight months earlier), we found measurable bupivacaine concentrations in the venous blood of some patients when the tourniquet cuff was still inflated, even though the exsanguination appeared adequate.<sup>8</sup> The three patients with the highest bupivacaine concentrations during the tourniquet inflation period (table 1) were asked to participate in a phlebographic examination to investigate possible venous leakage. In addition, three volunteers of normal build and arterial blood pressure (table 1) also were investigated. The build of the arm of the sub-

Within a few seconds of these first symptoms, convulsions occurred. No arterial blood pressure readings could be obtained. Venous blood was sampled with a  $P_{50}$  of 1.5, and thiopental (100 mg) and succinylcholine (100 mg) were given. The trachea was intubated and endotracheal intubation initiated. During approximately 15 min of mechanical ventilation, the patient was given 0.5 mg atropine after (given twice), 1.5 g rocuronium, and 1 mg suxamethonium. The total dose of sodium bicarbonate was 170 mEq, based on arterial blood gas analysis 10 min (pH 7.08,  $P_{50}$  100 mmHg, BE -14.4 mEq/L) and 25 min (pH 7.19,  $P_{50}$  45.8 mmHg, BE -11.1 mEq/L) after the last of convulsions. After the initial resuscitation the circulation was maintained with dopamine and intravenous infusions. The patient regained consciousness 30 min later, and the cardiovascular status returned to its preoperative level (arterial blood pressure 140/90 mmHg). After 40 min, analysis of arterial blood gas was normal.

25 and 40 min after the first observation of the toxic symptoms the arterial bupivacaine blood concentrations were 1.8 µg/ml, 0.9 µg/ml, and 0.8 µg/ml, respectively. Bupivacaine levels were determined by gas chromatography. The tourniquet, the cuff of which had remained inflated at 200 mmHg, was deflated after 30 min under (disinflation) (100 mmHg) anesthesia. One minute after the release, the bupivacaine concentration in the ipsilateral subcutaneous venous blood was 1.05 µg/ml, and five min later the concentration was 0.65 µg/ml. No convulsions or cardiac arrhythmias occurred at that time. The recovery was good and no signs of hypoxic organ damage were evident.

Two days later, phlebography of the same upper extremity was performed to visualize the arterial bupivacaine leakage. Information of the patient was obtained. The arm was exsanguinated with an Esmarch bandage in the same way as the IVRA with a cuff pressure of 200 mmHg (arterial pressure 180/110 mmHg). Eighty milliliters of contrast medium (Urografin® 360) were injected into a distal arterial vein in 20 s, and roentgenograms were taken every 10 s for 2 min, and then again after 5 min. Leakage of the contrast medium under the inflated cuff (the same as that seen on the patient two days earlier) was seen in the first roentgenogram (fig. 1) and thereafter. The arterial blood pressure during the phlebography remained at 170–180/100–110 mmHg.

CLINICAL REPORTS 96

Anesthesiology  
Vol. 58, No. 1, Jan 1983




FIG. 1. Phlebography of the arm after injection of contrast medium into a peripheral vein of the exsanguinated extremity. The tourniquet cuff pressure was 200 mmHg, and the same as when this position was given in IVRA with convulsions. Note the distal arterial vein on the proximal side of the cuff area (the right, curved arrow) and a continuous narrow contrast medium vein on the medial side of the biceps (the straight arrow).

#### RESULTS

In two of the patients, contrast medium was seen on the proximal side of the cuff as soon as 15 s after the injection. The filling of a short distance of some of the larger veins on the proximal side did not change during the 5-min examination, and no indications of open veins under the cuff could be seen. In the third patient (number 3) no contrast medium passed the tourniquet cuff (fig. 2). However, in two of the three volunteers (anesthesiologists), contrast medium leakage was observed at a cuff pressure of 250 mmHg. In the female volunteer some very narrow streaks of contrast medium were visible on the roentgenograms, from 15 s to 3 min, and the larger veins were filled on the proximal side of the cuff (fig. 3). The tourniquet cuff pressure appeared to be sufficient to block the arterial inflow, however, as the pale skin color remained unchanged during the study.

#### TABLE 1. Characteristics and Arterial Blood Pressure of the Patients and Volunteers during Previous IVRA and Present Phlebography of the Arm. Volume and Leakage of Contrast Medium Also Are Indicated

Patients	Sex	Age	Weight (kg)	Height (cm)	IVRA Pressure (mmHg)	Phlebography (mmHg)			Cuff pressure (mmHg)	Volume (ml)	Leakage (ml)	Bupivacaine concentration (µg/ml)
						Pre-injection	Mid-injection	Post-injection				
1	M	52	86	178	160/115	170/118	220/130	180/115	300	80 (0.00)*	+	0.80‡
2	M	41	94	186	120/80	140/118	160/118	130/110	250	80 (0.00)	+	0.80
3	M	29	71	173	120/80	140/118	160/118	140/110	250	80 (0.00)	-	0.81
Volunteers												
1	F	37	60	163		140/98	130/100		250	50	+	
2	M	39	55	182		130/90	140/100		250	50	+	
3	M	37	77	185		133/73	140/98		300	60	-	

\* Volume of 0.25% bupivacaine used for the preceding IVRA in the patients.  
† = leakage of contrast medium under the cuff; - = no leakage of contrast medium.  
‡ Concentration of bupivacaine in venous blood during the preceding IVRA.